

On the longitudinal structure function in the dipole model

MAREK NIEDZIELA^{(1),(2)}, MICHAŁ PRASZALOWICZ⁽¹⁾

⁽¹⁾M. Smoluchowski Institute of Physics, Jagiellonian University,
S. Łojasiewicza 11, 30-348 Kraków, Poland.

⁽²⁾Present address: Institute for Experimental Physics, University of Hamburg,
Luruper Chaussee 149, D-22761 Hamburg, Germany.

We compare new HERA data for the longitudinal structure function F_L with the predictions of different variants of the dipole model. In particular we show that the ratio F_L/F_2 is well described by the dipole models and is rather insensitive to the details of the fit. Fits to F_2 are performed with the help of geometrical scaling (GS). Using the property of GS we derive the bounds for F_L/F_2 both for the different versions of the dipole model and in the general case. Finally we briefly discuss how the higher Fock components of the photon wave function may affect these bounds.

PACS numbers: 13.85.Ni, 12.38.Lg

1. Introduction

Recently H1 [1] and ZEUS [2] Collaborations have published new data on the longitudinal structure function $F_L(x, Q^2)$ in deep inelastic ep scattering (DIS). The H1 analysis extends and improves previous data [3], which now cover kinematical range from $Q^2 = 1.5 \text{ GeV}^2/c^2$ and $x = 0.279 \times 10^{-4}$ up to $Q^2 = 800 \text{ GeV}^2/c^2$ and $x = 0.0322$. ZEUS data has been taken in much smaller region from $Q^2 = 9$ up to $Q^2 = 110 \text{ GeV}^2/c^2$ (see [4] for summary). In both data sets there is strong correlation between Q^2 and x values; for each Q^2 structure function F_L (and also F_2 that has been measured in the same kinematical points) has been measured over a limited x range, with small x 's concentrated around small values of Q^2 , see Fig. 1. Moreover, since F_L is difficult to extract experimentally, even recent improved data has still large errors.

Longitudinal structure function is of particular interest since, in the first approximation of the parton model, it is equal identically zero [5] (Callan-Gross relation) and therefore it is generated entirely by radiative corrections.

On the contrary, in the dipole model F_L is nonzero, albeit small. Indeed, Nachtmann and collaborators have shown that in the dipole model there exists a strict bound that [6, 7, 8]

$$F_L \leq g_{\max} \times F_2 = 0.27 \times F_2. \quad (1)$$

This result, hereafter referred to as an EMNS bound, is independent of the dipole-proton cross-section, and – strictly speaking – follows solely from the properties of the photon- $\bar{q}q$ wave function.




Fig. 1. H1 (blue circles) and ZEUS (red squares) data [1, 2] for F_2/Q^2 (left) and F_L/Q^2 (right) plotted as functions of Q^2 . Different points for one value of Q^2 correspond to different Bjorken x 's.

Using previous H1 data [3] the authors of Ref. [8] have shown that the bound (1) was almost saturated, which is difficult to realize in realistic dipole models. In this paper we are going to check if this conclusion remains still valid for the new data and what is the value of bound (1) for commonly used dipole-proton cross-sections. Similar analysis for the Golec-Biernat–Wüsthoff model [9, 10] has been already performed in Ref. [11]. To simplify the analysis we use here the property of geometrical scaling [12] (GS) which is to large precision exhibited by the DIS data up to relatively large Bjorken x 's [13]. We find that for realistic dipole-proton cross-sections the bound is indeed lower than (1) with $g \simeq 0.22$ (*i.e* 18.5% below the bound of Eq. (1)). In reality these bounds would be lowered if charm quark mass was included.

With present experimental accuracy we do not find any significant tension between F_L/F_2 data and the dipole model. Should such tension arise when new data appear, higher order corrections to the dipole model might resolve the issue. Therefore we discuss a possibility that corrections to the dipole model coming from higher Fock states in the virtual photon may change (1). Higher Fock components are needed *e.g.* in the dipole model [9] description of the diffractive data [10]. We show that this is possible only if there exists a bound for higher Fock components that is analogous to (1),

but with longitudinal contribution to F_2 that is significantly different than in the case of the $\bar{q}q$ state. Only explicit calculation of the $\bar{q}qg$ contribution to the virtual photon wave function might give here a definite answer. Such calculations have been recently carried out by various authors [14, 15, 16], however these results have not been so far applied to the phenomenological analysis of DIS. Although the calculation of $F_L^{\bar{q}qg}/F_2^{\bar{q}qg}$ with the help of Refs. [15, 16] might be probably possible, it is beyond the scope of the present paper.

The paper is organized as follows. In Sect. 2 we rederive the EMNS bound using geometrical scaling. Next, in Sect. 3, we fit two versions of the dipole model to the present data set for F_2 . To this end we also use the property of geometrical scaling. We then compare these fits with the data for F_L and discuss fit uncertainties. In Sect. 4 we calculate ratio F_L/F_2 for the afore mentioned fits and compare it with the data and with the EMNS bound. We do not find large tension between the data and model predictions. An influence of higher Fock states on the EMNS bound is discussed in Sect. 5. We conclude in Sect. 6.

2. Geometrical scaling and the EMNS bound

For three massless flavors DIS structure functions read [17]:

$$F_2(x, Q^2) = \frac{Q^2}{4\pi^3} \int dr^2 \left\{ |\psi_T(r, Q^2)|^2 + |\psi_L(r, Q^2)|^2 \right\} \sigma_{dp}(r^2),$$

$$F_L(x, Q^2) = \frac{Q^2}{4\pi^3} \int dr^2 |\psi_L(r, Q^2)|^2 \sigma_{dp}(r^2)$$

where photon wave functions take the following form

$$|\psi_T(r, Q^2)|^2 = \int_0^1 dz [z^2 + (1-z)^2] \bar{Q}^2 K_1^2(\bar{Q}r),$$

$$|\psi_L(r, Q^2)|^2 = 4 \int_0^1 dz z(1-z) \bar{Q}^2 K_0^2(\bar{Q}r). \quad (2)$$

Here K_i are modified Bessel functions and

$$\bar{Q}^2 = z(1-z)Q^2. \quad (3)$$

It is convenient to define functions $\Phi_{T,L}$

$$\Phi_{T,L}(u = rQ) = r^2 |\psi_{T,L}(r, Q^2)|^2 \quad (4)$$

that depend only on the combined variable $u = Qr$. Therefore

$$\begin{aligned} F_2(x, Q^2) &= \frac{Q^2}{2\pi^3} \int du \{ \Phi_T(u) + \Phi_L(u) \} \frac{\sigma_{dp}(u/Q)}{u}, \\ F_L(x, Q^2) &= \frac{Q^2}{2\pi^3} \int du \Phi_L(u) \frac{\sigma_{dp}(u/Q)}{u}. \end{aligned} \quad (5)$$

This parametrization is very convenient for the following reasons. First of all wave functions $\Phi_{T,L}(u)$ are universal and do not depend on external kinematical variables. Secondly, unlike functions $\psi_{T,L}(r, Q^2)$, they are everywhere regular in u . And finally, cross-section $\sigma_{dp}(u/Q)/u$ is a localized function of variable u that vanishes both for $u \rightarrow 0$ and $u \rightarrow \infty$.

If – as it is in the case of the GBW model – the dipole-proton cross-section exhibits geometrical scaling, *i.e.* $\sigma_{dp}(r) = \sigma_{dp}(rQ_s(x))$ then the integral

$$\int du \Phi_{T,L}(u) \frac{\sigma_{dp}(u/Q)}{u} = \int du \Phi_{T,L}(u) \frac{\sigma_{dp}(uQ_s/Q)}{u} = \text{function}(\tau) \quad (6)$$

depends only on a scaling variable

$$\tau = (Q/Q_s)^2. \quad (7)$$

Here Q_s^2 is a saturation scale

$$Q_s^2 = Q_0^2 \left(\frac{x}{x_0} \right)^{-\lambda}. \quad (8)$$

Now we can rederive the EMNS bound by considering the ratio

$$\frac{F_L(\tau)}{F_2(\tau)} = \frac{\int du \Phi_L(u) \sigma_{dp}(u/Q)/u}{\int du \{ \Phi_T(u) + \Phi_L(u) \} \sigma_{dp}(u/Q)/u} \quad (9)$$

and observing that function

$$g(u) = \frac{\Phi_L(u)}{\Phi_T(u) + \Phi_L(u)}, \quad (10)$$

which is plotted in Fig. 2, has a maximum $g_{\max} = 0.2714$ for $u = 2.591$. Therefore it follows that

$$\frac{F_L(x, Q^2)}{F_2(x, Q^2)} \leq g_{\max} = 0.27. \quad (11)$$




Fig. 2. Solid (red) line: function $g(u)$ defined in Eq. (10). Short dashed (brown) line: function $g(u)$ in the case of massive quarks for $\delta_f = m_f/Q = 0.5$ and long dashed (blue) line: for $\delta_f = m_f/Q = 1.06$, which corresponds to the charm quark mass $m_c = 1.3 \text{ GeV}/c$ and $Q^2 = 1.5 \text{ GeV}^2/c^2$.

The bound (11) has been derived for the case of massless quarks. While this is certainly a good approximation for three light flavors, given the fact that the lowest photon virtuality in the data set we use is $Q^2 = 1.5 \text{ GeV}^2/c^2$, the inclusion of charm mass effects is going to change (11). For a given flavor function g defined in Eq. (10) is no longer a function of scaling variable u only but in addition depends on the ratio $\delta_f^2 = m_f^2/Q^2$. For large Q^2 (*i.e.* small δ_f) $g(u, \delta_f) \rightarrow g(u)$. Moreover we have found numerically that everywhere in u we have

$$g(u, \delta_f) \leq g(u) \quad (12)$$

and the maxima $g_{\max}^{(f)}$ of $g(u, \delta_f)$ are decreasing with increasing δ_f , as illustrated in Fig. 2. This is consistent with the observation of Ref. [6] that $g_{\max}^{(f)}$ is a monotonically growing function of Q^2 . Therefore

$$0 \leq g_{\max}^{(f)}(Q^2) \leq g_{\max}. \quad (13)$$

This allows us to estimate the effect of the charm quark on the ratio:

$$\begin{aligned} \frac{F_L^{(\text{light}+c)}}{F_2^{(\text{light}+c)}} &= \frac{F_L + F_L^{(c)}}{F_2 + F_2^{(c)}} = \frac{F_L/F_2 + F_L^{(c)}/F_2^{(c)} F_2^{(c)}/F_2}{1 + F_2^{(c)}/F_2} \\ &\leq g_{\max} \frac{1 + g_{\max}^{(c)}/g_{\max} F_2^{(c)}/F_2}{1 + F_2^{(c)}/F_2} \leq g_{\max} \end{aligned} \quad (14)$$

where the last inequality follows from (13). Note that $F_{2,L}$ without any superscript refers to the structure function with light flavors only and that dependence on Q^2 has been suppressed. We see therefore, in agreement with

x_{\max}	$\sigma_0[1/\text{GeV}^2]$	λ	x_0	χ^2/dof
none	23.68	0.389	0.010497	1.18
0.01	27.11	0.353	0.007786	0.87
0.005	29.33	0.333	0.006435	0.79
0.0005	38.37	0.253	0.003090	0.70

Table 1. Parameters of the GBW model fitted to F_2 H1 data up to x_{\max} .

Ref. [8], that inclusion of charm lowers the bound on $F_L^{(\text{light}+c)}/F_2^{(\text{light}+c)}$ in proportion that depends on $F_2^{(c)}/F_2$. For $m_c = 1.3 \text{ GeV}/c$ and for the lowest Q^2 in the present data set we get numerically $g_{\max}^{(c)} \approx 0.05$ (see Fig. (2)), which gives $g_{\max}^{(c)}/g_{\max} \approx 0.19$. We do not know what is the fraction of charm in the present data sample, however taking a typical value of $F_2^{(c)}/F_2 \sim 25\%$, we get that $g_{\max}^{(\text{light}+c)} \sim 0.23$. This means that bound (11) is lowered for the lowest Q^2 bin by approximately 18% and is approaching (11) for higher Q^2 . In the following we will ignore charm contribution and stick to the bound (11) coming back to this point in Sect. 4.

3. Dipole models and geometrical scaling

In order to check how far the bound (11) overestimates actual predictions of the dipole model with realistic dipole-proton cross-section, we are going to compute ratio (11) for a given σ_{dp} in terms of scaling variable τ . We will see that for commonly used parametrizations of σ_{dp} , ratio F_L/F_2 is a slowly varying function of τ with a maximum equal approximately 0.216 – 0.224, which only slightly depends on the parametrization actually used. To this end we have decided to perform our own fits to the F_2 data over the restricted kinematical range where the longitudinal structure function F_L has been measured by H1. The reason for this is threefold. Firstly, new data is of much better quality than the previous data to which the dipole model parameters have been fitted. Secondly, we do not aim at a global fit, but rather at a fit which covers only the points where F_L has been measured as well. Therefore fit parameters – as we shall see in the following – will be different from the ones obtained in the global fits. And finally, we have decided to fit the data with the help of geometrical scaling – a procedure not used so far in the fits to the DIS data.

Fitting dipole models to the data becomes very easy when F_2 depends only upon single scaling variable τ . This happens because points corresponding to one particular value of Q^2 but different x 's (see "stacks" in Fig. 1) correspond to different values of τ and are therefore shifted horizontally – if plotted in terms of τ – by values that are different for different

x 's. As a consequence dipole model predictions fall on a universal curve (up to an overall normalization σ_0), and data fitting consists in changing σ_0 and the parameters defining scaling variable τ , *i.e.* x_0 and λ . By varying these three parameters one forces experimental points to fall on theoretical prediction, rather than by changing theoretical predictions one is trying to reproduce experimental points. Therefore this method is very efficient, as it does not require time consuming recalculations of the theoretical curve.

Even in the case of dipole-proton cross-sections that violate GS by explicit (albeit weak) dependence on x , like in the model of Iancu, Itakura and Munier [18], it is still possible to make a GS-like fit by keeping x at some fixed average value x_{ave} and then study the uncertainty of theoretical predictions by changing x over the range that is covered by experimental data. We shall come back to this point later.

Let us first consider the simplest version of the dipole model, namely the GBW parametrization [9], which – up to an overall normalization σ_0 – takes the following form in terms of scaling variable τ

$$\frac{\sigma_{\text{dp}}^{\text{GBW}}(u, \tau)}{\sigma_0} = 1 - \exp\left(-\frac{u^2}{4\tau}\right). \quad (15)$$

Plugging (15) into Eqs.(5) gives unnormalized theoretical prediction for the structure functions divided by Q^2 , which will be denoted in the following by small characters $f_{2,L}(\tau)$. Experimental data are tabulated in a set of discrete points $\{Q_i^2, x_i\}$, and we fit three free parameters of the model, σ_0, x_0 and λ , by transforming experimental entries in the following way:

$$F_2(x_i, Q_i^2) \rightarrow \frac{1}{Q^2 \sigma_0} F_2(x_i, Q_i^2) = f_2\left(\tau_i = \frac{Q^2}{Q_s^2(x_i)}\right) \quad (16)$$

and demanding that they are equal to the theoretical prediction at the pertinent value of scaling variable τ_i with Q_s given by Eq.(8). The results are shown in Table 1. Since GS is supposed to work the best for small values of Bjorken x 's we have performed a number of fits restricting the allowed x region up to a maximal value denoted as x_{max} . We see that even without any cut on the maximal value of x , *i.e.* for x as large as 0.0322 (the highest x in the analyzed data set) χ^2 of the fit is quite reasonable. By restricting analyzed data to the smaller and smaller range of Bjorken x 's χ^2 is getting smaller, but also model parameters vary substantially. Parameter σ_0 is much smaller than in the original fit of Ref. [9] $\sigma_0^{\text{GBW}} = 23 \text{ mb} = 59 \text{ GeV}^{-2}$. Exponent λ approaches the value of Ref. [9] $\lambda^{\text{GBW}} = 0.288$ only for small x_{max} (note that maximal x in Ref. [9] was equal to 0.01, whereas the lowest $x = 6 \times 10^{-6}$ was much below the minimal x of present analysis), otherwise being consistent with model-independent analysis of Ref. [13]. The results

of the fits, together with the original parametrization of Ref. [9] are plotted in Fig. 3. One can see rather good agreement of fits from Table 1 with the data, and – also quite importantly – good quality of GS of the data.




Fig. 3. H1 (blue circles) and ZEUS (red squares) data [1, 2] for F_2/Q^2 plotted as a function of scaling variable τ for different values of fit parameters corresponding to the GBW model given in Table 1. Fit parameters in the first panel correspond to the original fit of Ref. [9] with no charm data included.

Finally in Fig. 4 we plot data for F_L/Q^2 as a function of τ together with theoretical parametrization of Ref. [9] and the present fits corresponding to Tab. 1. We can see that due to still large experimental errors of F_L all parametrizations, although different, describe well the data.

As a second example let us consider a dipole model by Iancu, Itakura and Munier (IIM) [18], where the dipole-proton cross-section is defined in terms of two functions

$$A_1(u, \tau, x) = A_0 \left(\frac{u^2}{4\tau} \right)^{\gamma + \frac{\ln(4\tau/u^2)}{2\kappa\lambda \ln(1/x)}},$$

$$A_2(u, \tau) = 1 - \exp(-a \ln^2(bu/\sqrt{\tau})) \quad (17)$$




Fig. 4. H1 (blue circles) and ZEUS (red squares) data [1, 2] for F_L/Q^2 plotted as a function of scaling variable τ . Solid curves correspond to the GBW model fits of Table 1.


and

$$\frac{\sigma_{\text{dp}}^{\text{IIM}}(u, \tau, x)}{\sigma_0} = \begin{cases} A_1(u, \tau, x) & \text{for } u^2 < 4\tau \\ A_2(u, \tau) & \text{for } 4\tau \leq u^2 \end{cases}. \quad (18)$$

Here $\gamma = 0.63$ is related to the anomalous dimension of the forward scattering amplitude in the BFKL formalism, while $\kappa = 9.9$ corresponds to the diffusion coefficient. Parameters a and b are determined uniquely by gluing A_1 and A_2 and their derivatives at $u^2 = 4\tau$. Parameter A_0 is in principle free, however, as it was shown in Ref. [18] the best χ^2 was obtained for $A_0 = 0.7$ and for the purpose of the present work we will keep it fixed at this value. Therefore the only free parameters are, as in the case of the GBW model, an overall normalization σ_0 and two parameters of the saturation scale: λ and x_0 .

However, there are two important differences between IIM and GBW parametrizations. First of all, for small value of u amplitude A_1 exhibits explicit violation of GS, since it depends both on τ and x . For the purpose of the present work we will keep x entering A_1 fixed at the average value x_{ave} calculated for the interval where the fit is performed. The accuracy of

x_{\max}	x_{ave}	$\sigma_0[1/\text{GeV}^2]$	λ	x_0	χ^2/dof
none	0.00359	20.22	0.597	0.002553	1.76
0.01	0.00182	21.50	0.583	0.002140	1.57
0.005	0.00121	25.56	0.531	0.001392	1.31
0.0005	0.00022	34.30	0.389	0.000645	0.75

Table 2. Parameters of the IIM [18] model fitted to F_2 H1 data up to x_{\max} .Fig. 5. H1 (blue circles) and ZEUS (red squares) data [1, 2] for F_2/Q^2 plotted as a function of scaling variable τ for different values of IIM model fit parameters given in Table 2.

this procedure is then checked by putting in (18) x equal to the maximal and minimal value of x in a given interval. It will turn out that the structure functions are sensitive to this variation of x at the level of a few percent, however the ratio F_L/F_2 is almost independent. Next difference appears due to the dependence of A_1 on λ . To solve this issue we have adopted an iterative procedure, consisting in fixing λ at some initial value, then performing the fit and plugging in the fitted value of λ to A_1 and performing the fit again. Usually after four, five steps satisfactory convergence has been achieved. The results are given in Table 2.

By inspecting Table 2 we see that the quality of fits is worse than in the




Fig. 6. $F_2(\tau)/Q^2$ for the parameters from the third row of Table 2 with x in A_1 from Eq. (17) replaced by x_{\min} (upper curve), x_{ave} (solid middle curve) and x_{\max} (lower curve).

case of the GBW model. This is in contrast with the original fits of Ref. [18] which, however, were performed over the data set covering much lower x 's than in our present analysis. One should also note that the errors of the old data sets are bigger than the ones of the present data. This is also the reason why fit parameters are different than in our case. For illustration purposes we have plotted in Fig. 5 $F_2(\tau)/Q^2$ for the original IIM parameters and for three choices of x_{\max} from Table 2. Magnifying first plot in Fig. 5 one could see that for $\tau > 10$ the original IIM curve missed the experimental points, which have rather small errors.

In order to check sensitivity of the IIM fits to the fact that x dependent piece of A_1 amplitude (17) has been replaced by a constant value x_{ave} , we plot in Fig. 6 $F_2(\tau)/Q^2$ for the parameters from the third row of Table 2 with x in A_1 replaced by x_{\min} (upper curve) and x_{\max} (lower curve). For better resolution the plot is restricted to $\tau < 5$. We can see that theoretical uncertainty introduced by this procedure is in fact much smaller than the experimental errors.

Finally in Fig. 7 we plot F_L/Q^2 as a function of τ for three sets of parameters from Table 2 and for the original set of parameters from Ref. [18]. One can see that all curves describe the data reasonably well due to the large error bars of F_L .

4. EMNS bound for dipole models

Now we are able to compare the EMNS bound with F_L/F_2 ratio calculated in the dipole model for realistic dipole-proton cross-sections σ_{dp} discussed in the previous Section. One should note that for the GBW dipole model F_L/F_2 does not depend on the values of x_0 , λ and σ_0 . It is not the




Fig. 7. H1 (blue circles) and ZEUS (red squares) data [1, 2] for F_L/Q^2 plotted as a function of scaling variable τ . Solid curves correspond to the IIM model fits of Table 2.




Fig. 8. Ratios F_L/F_2 as functions of τ for the GBW dipole model and for the IIM model with parameters corresponding to the fit with $x < 0.005$. The scale of the plot has been enlarged in order to make small differences between the two curves visible.

case for the IIM dipole model, but we have checked explicitly that for all parametrizations of Table 2 the differences in F_L/F_2 are negligible. Therefore in Fig. 8 we plot ratios F_L/F_2 as functions of τ for the GBW dipole

model	τ_{\max}	$(F_L/F_2)_{\max}$
GBW any	1.165	0.224
IIM all x	1.417	0.217
IIM $x < 0.01$	1.411	0.216
IIM $x < 0.005$	1.413	0.216
IIM $x < 0.0005$	1.418	0.217

Table 3. Maxima of F_L/F_2 for different fits to F_2 .

model and for the IIM model with parameters corresponding to the fit with $x < 0.005$. We see that ratios F_L/F_2 are in fact almost model-independent. This is further confirmed in Table 3 where we collect the maximal value of F_L/F_2 for the GBW and IIM parametrizations.

One should note that each curve in Fig. 8 corresponds to a different definition of scaling variable τ , so one cannot superimpose experimental data on that plot. This is done in Fig. 9 where we plot F_L/F_2 for the unrestricted fits of the GBW and IIM models corresponding to the first rows of Table 1 and 2 respectively. The errors of the ratio have been calculated neglecting correlation between errors of F_2 and F_L :

$$\Delta \left(\frac{F_L}{F_2} \right) = \frac{F_L}{F_2} \sqrt{\left(\frac{\Delta F_L}{F_L} \right)^2 + \left(\frac{\Delta F_2}{F_2} \right)^2} \quad (19)$$

This procedure overestimates the errors, however, given the fact that $F_L < 0.27F_2$ and that experimentally absolute errors of F_L are 2 – 10 times larger than ΔF_2 , the error of the ratio (19) is determined to very high precision by ΔF_L alone. We shall come back to this point later.




Fig. 9. Ratios F_L/F_2 plotted as functions of scaling variable τ . Straight line corresponds to the EMNS bound. Solid line in the left panel corresponds to the unrestricted GBW fit of Tab. 1 (first row), and in the right panel to the unrestricted IIM fit of Tab. 2 (first row).

We can see from Figs. 8 and 9 that for realistic σ_{dp} theoretical predictions lie below the EMNS bound. Indeed, we see that the maximum of F_L/F_2 is of the order $0.216 - 0.224$ and only slightly varies from fit to fit. This is illustrated in Table 3. A question arises whether data points – which for some values of τ exceed the EMNS bound – are indeed, as suggested by the authors of Ref. [8], saturating bound (11), being as a consequence incompatible with the dipole model. To this end we have simply calculated χ^2 of F_L/F_2 for the unrestricted GBW fit and obtained very good result: 0.7425. To check whether this value is affected by the fact that we have not taken into account correlations between errors of F_L and F_2 , we have recalculated χ^2 neglecting F_2 errors, which gives χ^2 that changes by less than 1%. Therefore indeed, as already mentioned above, χ^2 value is totally driven by the large errors of F_L .

It is interesting to check whether an overall shift of the dipole model prediction for F_L/F_2 would improve agreement with the data. In this way we shall have a quantitative measure of the quality of the dipole model prediction for the ratio F_L/F_2 and also an indication how much room is there for the higher order corrections that we are going to discuss in Sect. 5. To this end we use the GBW fits allowing for arbitrary normalization of the ratio

$$\frac{F_L}{F_2} \rightarrow \mathcal{N} \frac{F_L}{F_2} \quad (20)$$

and calculate $\chi^2/\text{d.o.f.}$ (assuming one degree of freedom, namely \mathcal{N}) as a function of \mathcal{N} . This is illustrated in Fig. 10 for two GBW fits of Table 1: unrestricted x and $x < 0.0005$. We see that depending on the fit the data prefers \mathcal{N} slightly smaller (unrestricted x) or slightly larger ($x < 0.0005$) than 1. These are negligible changes and therefore at this stage we conclude that the data for the ratio F_L/F_2 is compatible with the dipole model. One should perhaps remind here again, that this ratio is quite stable, even for fits to F_2 that are visibly different.

One can also see that lowering \mathcal{N} by 20%, which would be required by taking into account charm mass effects (*c.f.* discussion at the end of Sect. 2) increases χ^2 , but only by a small amount. Of course the detailed study of charm mass effects would require to go beyond GS used in this analysis – which in turn would result not only in the change of \mathcal{N} , but also in the change of the shape of F_L/F_2 – but no drastic difference to the present analysis should be expected.

5. EMNS bound and higher Fock states

In the dipole model the virtual photon dissociates into a $\bar{q}q$ pair which subsequently interacts with the proton target. However, it is clear that




Fig. 10. Change of χ^2 of $\mathcal{N} \times F_L/F_2$ as a function of \mathcal{N} for the GBW fits from Table 1 for unrestricted x (left) and $x < 0.0005$ (right).

higher Fock states have to contribute as well, similarly to the diffractive DIS where the next Fock component, namely the $\bar{q}qg$ state, is dominant at small β [10]. Full calculations of the $\bar{q}qg$ component of the photon wave function have recently appeared in the literature [14, 15, 16], however they have not been yet applied to the phenomenological analysis of DIS.

Structure functions in the dipole model are given as an expansion

$$F_{2,L}(x, Q^2) = F_{2,L}^{(\bar{q}q)}(x, Q^2) + F_{2,L}^{(\bar{q}qg)}(x, Q^2) + \dots \quad (21)$$

The EMNS bound (11) derived in Sect. 2 is in fact valid only for the first component of (21)

$$G = \frac{F_L^{(\bar{q}q)}(x, Q^2)}{F_2^{(\bar{q}q)}(x, Q^2)} < g_{\max} = 0.27. \quad (22)$$

One should note, however, that at this order of perturbative expansion the loop corrections to the leading order Fock component may change the value of the bound (22). Similarly for the next Fock component we would have:

$$H = \frac{F_L^{(\bar{q}qg)}(x, Q^2)}{F_2^{(\bar{q}qg)}(x, Q^2)} < h_{\max} \quad (23)$$

where

$$0 \leq h_{\max} \leq 1. \quad (24)$$

Up to this order one can derive the *modified* EMNS bound:

$$\begin{aligned}
\frac{F_L(x, Q^2)}{F_2(x, Q^2)} &= \frac{F_L^{(\bar{q}q)}(x, Q^2) + F_L^{(\bar{q}qg)}(x, Q^2)}{F_2^{(\bar{q}q)}(x, Q^2) + F_2^{(\bar{q}qg)}(x, Q^2)} \\
&< \frac{g_{\max} F_2^{(\bar{q}q)}(x, Q^2) + h_{\max} F_2^{(\bar{q}qg)}(x, Q^2)}{F_2^{(\bar{q}q)}(x, Q^2) + F_2^{(\bar{q}qg)}(x, Q^2)} \\
&= g_{\max} \frac{1 + \delta \varepsilon(x, Q^2)}{1 + \varepsilon(x, Q^2)}
\end{aligned} \tag{25}$$

where

$$\delta = \frac{h_{\max}}{g_{\max}}, \quad 0 \leq \delta \leq \frac{1}{g_{\max}} \simeq 3.7$$

and

$$\varepsilon(x, Q^2) = \frac{F_2^{(\bar{q}qg)}(x, Q^2)}{F_2^{(\bar{q}q)}(x, Q^2)}. \tag{26}$$

Since the dipole model with $\bar{q}q$ component only describes F_2 rather well over the wide kinematical range, we do not expect ε to be large. For the purpose of the present analysis we assume that ε does not exceed 20%.




Fig. 11. Modification factor f of Eq. (27) as a function of ϵ and δ . Light green plane corresponds to $f = 1$.

In order to change the value of the EMNS bound we need the *modification factor*

$$f = \frac{1 + \varepsilon \delta}{1 + \varepsilon} \tag{27}$$

to be significantly different from 1. In Fig. 11 we plot $f(\varepsilon, \delta)$ for $0 \leq \varepsilon \leq 0.2$ and $0 \leq \delta \leq 1/g_{\max}$. We see from Fig. 11 and from Eq. (27) that $f = 1$ for $\delta = 1$ and that f gets smaller than 1 if $\delta < 1$ and $f > 1$ for $\delta > 1$, and that the difference $|f - 1|$ is growing with ε . From Fig. 11 one may conclude that the modification of the EMNS bound by more than $\pm 20\%$ might be rather difficult. Most probably higher Fock components would modify (11) by less than 10%, but to quantify this statement one needs to calculate explicitly h_{\max} which is beyond the scope of the present paper. One should note at this point that in this case an inclusion of mass corrections due to the charm and possibly bottom quarks should be included, as these corrections would be of the same order or even larger than a contribution from the higher Fock components.

To conclude this Section let us only remark that ε depends on kinematical variables, so one cannot exclude *a priori* a situation that there exists a kinematical corner where a correction to (11) is of importance.

6. Conclusions

Dipole model offers effective and intuitive description of deep inelastic scattering which goes beyond leading twist approximation. In this paper we have used different variants of the dipole model, *i.e.* two different forms of photon-proton cross-section fitted to F_2 data over different ranges of Bjorken x 's. We have decided to perform our new fits in the kinematical range where both F_2 and F_L have been measured. This kinematical region does not extend to very small x 's as it is in the case of the recently published combined HERA data [19] and therefore fitted parameters are different than the ones of global fits. For the same reason the IIM model [18] that has been specifically devised for low x region gives larger χ^2 than the simplest version of the the GBW model [9], which quite satisfactorily describes F_2 .

In order to fit model parameters we have used the property of geometrical scaling, which boils down to the fact that data points of the same Q^2 but different x 's disperse when plotted in terms of scaling variable τ and fall on one line (compare Figs. 1 and *e.g.* 3). We have found that GS is present both in the case of F_2 and F_L as well. One should, however, take this property with care for the present set of data, since for each value of Q^2 only a few points of different Bjorken x 's have been measured. This $x - Q^2$ correlation of the DIS data, particularly pronounced in the present case of F_L , is a common obstacle in deriving firm conclusions on the quality of GS. In this context it is also worth mentioning that there exist different sources of GS. The first one is related to the genuine property of the initial state of the proton target described theoretically by the Color Glass Condensate formalism [20] and nonlinearities in the parton evolution [21], which are of

importance at small values of Bjorken x . The other one is a property of the linear parton evolution equations [22], which build GS at larger Q^2 and at not necessarily very small x 's.

We have next compared dipole model predictions – with parameters fixed by fits to F_2 – with F_L data. Here, due to large experimental errors, agreement is quite good. Next we have studied ratio of the structure functions $G = F_L/F_2$ for which in the dipole model there exists a strict bound $g_{\max} = 0.27$ (1) derived in Ref. [8]. We have rederived (1) with the help of geometrical scaling. We have also shown that in the dipole models discussed above $g_{\max}^{\text{dp}} \approx 0.216 - 0.224$ which is approximately 18.5% below the EMNS bound. Different fits give very similar ratio g , which is only residually dependent on the values of fit parameters (this concerns only the IIM dipole model, since ratio F_L/F_2 is parameter independent in the GBW case). Comparing $G = F_L/F_2$ with the data we have established that the GBW model reproduces G with high precision. We do not see any tension between the data and the dipole model as far as ratio G is concerned, even if charm mass effects, which lower the EMNS bound, are taken into account.

Dipole model and the EMNS bound discussed so far rely on the first Fock component of the photon wave function. Including higher Fock components, like a $\bar{q}qg$ state, might in principle change theoretical prediction for g_{\max} . For this to happen, longitudinal part of the $\bar{q}qg$ state compared to the transverse one has to be significantly different than in the $\bar{q}q$ case.

Acknowledgments

The authors want to thank Aharon Levy for bringing to their attention new data on F_L and for an access to this data prior to publication. M.P. thanks Otto Nachtmann for a conversation that started this project and for remarks on the final version of the manuscript. We thank Guillaume Beuf for bringing to our attention Refs. [15, 16]. The research of M.P. has been supported by the Polish NCN grants 2011/01/B/ST2/00492 and 2014/13/B/ST2/02486.

REFERENCES

- [1] V. Andreev *et al.* [H1 Collaboration], Eur. Phys. J. C **74** (2014) 2814 [arXiv:1312.4821 [hep-ex]].
- [2] H. Abramowicz *et al.* [ZEUS Collaboration], arXiv:1404.6376 [hep-ex].
- [3] F. D. Aaron *et al.* [H1 Collaboration], Phys. Lett. B **665** (2008) 139 [arXiv:0805.2809 [hep-ex]] and Eur. Phys. J. C **71** (2011) 1579 [arXiv:1012.4355 [hep-ex]].
- [4] A. Levy, talk at QCD Moriond 2014, arXiv:1405.3753 [hep-ex].
- [5] C. G. Callan, Jr. and D. J. Gross, Phys. Rev. Lett. **22** (1969) 156.
- [6] C. Ewerz and O. Nachtmann, Phys. Lett. B **648** (2007) 279 [hep-ph/0611076].
- [7] C. Ewerz, A. von Manteuffel and O. Nachtmann, Phys. Rev. D **77** (2008) 074022 [arXiv:0708.3455 [hep-ph]].
- [8] C. Ewerz, A. von Manteuffel, O. Nachtmann and A. Schöning, Phys. Lett. B **720** (2013) 181 [arXiv:1201.6296 [hep-ph]].
- [9] K.J. Golec-Biernat and M. Wüsthoff, Phys. Rev. D **59** (1998) 014017 [hep-ph/9807513].
- [10] K.J. Golec-Biernat and M. Wüsthoff, Phys. Rev. D **60** (1999) 114023 [hep-ph/9903358].
- [11] C. Ewerz and O. Nachtmann, Annals Phys. **322** (2007) 1670 [hep-ph/0604087].
- [12] A.M. Stasto, K.J. Golec-Biernat, J. Kwiecinski, Phys. Rev. Lett. **86** (2001) 596.
- [13] M. Praszalowicz and T. Stebel, JHEP **1303** (2013) 090 [arXiv:1211.5305 [hep-ph]].
- [14] I. Balitsky and G. A. Chirilli, Phys. Rev. D **83** (2011) 031502 [arXiv:1009.4729 [hep-ph]] and Phys. Rev. D **87** (2013) 1, 014013 [arXiv:1207.3844 [hep-ph]].
- [15] G. Beuf, Phys. Rev. D **85** (2012) 034039 [arXiv:1112.4501 [hep-ph]].
- [16] R. Boussarie, A. V. Grabovsky, L. Szymanowski and S. Wallon, JHEP **1409** (2014) 026 [arXiv:1405.7676 [hep-ph]].
- [17] See *e.g.* Chapter 9 in V. Baronne and E. Predazzi, *High-Energy Particle Diffraction*, Springer 2002.
- [18] E. Iancu, K. Itakura and S. Munier, Phys. Lett. B **590** (2004) 199 [hep-ph/0310338].
- [19] F. D. Aaron *et al.* [H1 and ZEUS Collaboration], JHEP **1001** (2010) 109 [arXiv:0911.0884 [hep-ex]].
- [20] L. D. McLerran, and R. Venugopalan, Phys. Rev. **D49**, 2233 (1994), Phys. Rev. **D49**, 3352 (1994), and Phys. Rev. **D50**, 2225 (1994).
- [21] S. Munier and R. B. Peschanski, Phys. Rev. Lett. **91** (2003) 232001 [hep-ph/0309177], and Phys. Rev. D **69** (2004) 034008 [hep-ph/0310357].
- [22] F. Caola and S. Forte, Phys. Rev. Lett. **101** (2008) 022001 [arXiv:0802.1878 [hep-ph]].